

Patterns in Plant Leaf Flammability of Woodlands and Forests of Eastern Australia



Daniel William Krix

Thesis submitted for the degree of Doctor of Philosophy

February 2020

Supervisors: Dr Brad R. Murray and Dr Megan L. Phillips

Certificate of Original Authorship

I, Daniel Krix declare that this thesis, is submitted in fulfilment of the requirements for the award of Doctor of Philosophy, in the School of Life Sciences at the University of Technology Sydney.

This thesis is wholly my own work unless otherwise reference or acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

This document has not been submitted for qualifications at any other academic institution.

This research is supported by an Australian Government Research Training Program.

Production Note:

Signature: Signature removed prior to publication.

Date: 10/08/19

Acknowledgments

Firstly, I'd like to thank my supervisor Brad Murray, who took me on for my Honours project, and worked hard to secure a scholarship without which I'd not have been able to continue on to work on this thesis. He's given me the space to develop my own ideas and managed to pull me out of my 'rabbit holes', many, many times, and sometimes with much resistance on my part! His enthusiasm and kind support have been absolutely invaluable through all phases of this project. Also, it was he who first suggested I try R; my getting to grips with programming since then has been unbelievably rewarding. Thanks also to my co-supervisor Megan Phillips, and to Leigh Martin, who have both provided helpful comments on various aspects of this work.

The lab staff at UTS were also incredibly helpful, particularly Gemma Armstrong, who modified the lab for the leaf burning experiments, and let me continue using the muffle furnace in spite of my causing several heat-induced breakdowns due to its use in an application it was patently not designed for. My parents, Sharyn and Michael provided countless cups of tea and space to sort through samples in the Blue Mountains, along with storage for bags of soil and various bits of equipment from my many failed ideas. Also, thanks to the Sunday Coffee Boys, Luke Blattman, Guy Dernee, Hayden Fryer and Bob Lloyd, who listened to me bang on about my project so many times.

I'll be forever grateful to my wife Caroline, who first suggested I find a new career path, and who has supported me in this crazy endeavour for nearly a decade now. She has given me the time to follow this road out as far as it goes, and it's proven to be more satisfying than I'd ever imagined. Finally, my daughter Charlotte, and the little man who's

almost here, who let me see the world with new eyes, and desire to make it better for them,
in whatever small way I might.

Table of Contents

Certificate of Original Authorship	II
Acknowledgments.....	III
Table of Contents.....	V
List of Figures	IX
List of Tables	XV
List of Abbreviations in this Thesis.....	XVI
Abstract.....	XVIII
Chapter 1: General Introduction.....	1
1.1 Wildfire and the flammability of plant species	1
1.2 Leaf flammability attributes and leaf traits	5
1.3 Research novelty and significance	8
1.4 Study region	9
1.5 Thesis aims	12
1.6 Chapter structure and descriptions	12
Chapter 2: Landscape variation in plant leaf flammability is driven by leaf traits responding to environmental gradients	14
2.1 Introduction.....	14
2.2 Methods	17
2.2.1 Study region.....	17
2.2.2 Study sites and species	19

2.2.3 Leaf collection.....	20
2.2.4 Leaf trait and flammability measurements	21
2.2.5 Statistical analyses.....	23
2.3 Results	26
2.3.1 Leaf traits in relation to habitat and stratum.....	26
2.3.2 Leaf flammability attributes in relation to habitat and stratum	26
2.3.3 Relationships between leaf flammability attributes and leaf traits.....	29
2.4 Discussion	33
Chapter 3: Relationships among leaf flammability attributes and identifying low-leaf-	
flammability species at the wildland-urban interface	38
3.1 Introduction.....	38
3.2 Methods	41
3.2.1 Study species and leaf collection	41
3.2.2 Leaf flammability attributes	42
3.2.3 Measurements of leaf ignitibility, sustainability and combustibility	43
3.2.4 Measurement of leaf traits.....	45
3.2.5 Relationships among flammability attributes	45
3.2.6 Calculation of an Overall Leaf Flammability Index (OLFI)	46
3.2.7 Relationships between leaf flammability and leaf traits	47
3.3 Results	47
3.3.1 Relationships between ignitibility attributes	47

3.3.2 Relationships between sustainability attributes	48
3.3.3 Relationships between ignitibility, sustainability and combustibility attributes ..	48
3.3.4 Identification of low-leaf-flammability species	50
3.3.5 Relationships between leaf functional traits and low leaf flammability	51
3.4 Discussion	54
Chapter 4: Do plant leaves become more flammable with increasing radiant heat?	59
4.1 Introduction.....	59
4.2 Methods	61
4.2.1 Study region and species	61
4.2.2 Leaf collection and measurement of flammability attributes	61
4.2.3 Statistical analysis.....	64
4.3 Results	66
4.3.1 Responses of flammability attributes to increasing radiant energy	66
4.3.2 Evidence for convergence among species in their flammability attributes at higher radiant energy	66
4.4 Discussion	71
Chapter 5: Predicting changes in plant leaf flammability in response to increasing radiant energy: the importance of leaf functional traits.....	74
5.1 Introduction.....	74
5.2 Methods	75
5.2.1 Study species	75

5.2.2 Measurement of leaf traits and flammability attributes	76
5.2.3 Predictive modelling of TTI and ID	78
5.2.4 Predictive modelling of leaf flaming behaviour	80
5.2.5 Determining the importance of including leaf traits in predictive modelling	82
5.2.6 Model validation.....	85
5.3 Results	89
5.3.1 The importance of including leaf traits in predictive modelling	89
5.3.2 Comparison of prediction error in TTI between the predictive and validation datasets	95
5.3.3 Comparison of prediction error in ID between the predictive and validation datasets	98
5.3.4 Comparison of classification accuracy of leaf flaming between the predictive and validation datasets	101
5.4 Discussion.....	103
Chapter 6: General Discussion	107
6.1 Were the aims of this thesis met?	107
6.2 Future research directions	112
6.3 Conclusion	113
Thesis appendices	114
References	144

List of Figures

Fig. 1.1 (following page) Map of the location of the study sites, with the sites broad location shown in (a), their situation within the Blue Mountains (b), and the topography and waterways of the sites in (c). The red square delineates the area shown in (c). Exposed sites (woodland vegetation) and sheltered sites (closed forest vegetation) are indicated by green and red circles respectively. The altitude of the sites varied from 200 m to 470 m above sea level.	10
Fig. 2.1 Typical exposed woodland habitat on ridge tops (a) and sheltered forest habitat (b), in the study region. Despite large differences in abiotic conditions, these habitats can often be found within short distances of each other (< 500 m).	18
Fig. 2.2 (following page) Bean plots of leaf traits (a) LA, (b) LMA and (c) FMC by habitat and stratum. Blue beans are sheltered habitat, and red beans are exposed habitat. Darker beans represent overstorey species and lighter beans represent understorey species. The shaded areas of the beans represent the estimated density of the distribution with short black lines in the shaded areas of the beans showing the observations. Long black lines are means of observations for a given group.	27
Fig 2.3 (following page) Bean plots of flammability attributes (a) ignitibility, (b) sustainability and (c) combustibility by habitat and stratum. Blue beans are sheltered habitat, and red beans are exposed habitat. Darker beans represent overstorey species and lighter beans represent understorey species. The shaded areas of the beans represent the estimated density of the distribution with short black lines in the shaded areas of the beans showing the observations. Long black lines are means of observations for a given group.	29
Fig. 2.4 (previous page) Scatter plots of partial regressions of flammability attributes (y-axes) against leaf traits (x-axes). Blue rhombuses represent sheltered species, red triangles represent exposed species. Line of best fit from GLM model shown as an orange line (identical to best fitting PGLS model). The grey shaded area represents the 95% confidence interval of the mean.	33
Fig. 3.1 Temporal pathway of typical plant leaf combustion, here showing values from a single sample of <i>Corymbia gummifera</i> . Length of time for each flammability attribute is represented in the	

top of the figure (TTS, time to smoking; TTI, time to incandescence; TTF, time to flaming; SD, smoking duration; ID, incandescence duration; FD, flaming duration). The grey bar represents smoking, the orange bar represents incandescence, and the red bar represents flaming. Vertical dotted lines delineate physical combustion processes. 43

Fig. 3.2 (a) Correlations between pairs of flammability attributes. Both x and y axes are time measurements (s). Red circles indicate significant relationships. (b) Boxplots of the distribution of species' values for each of the flammability attributes (TTS, time to smoking; TTI, time to incandescence; TTF, time to flaming; SD, smoking duration; ID, incandescence duration; FD, flaming duration). 49

Fig. 3.3 Relationships between combustibility (flaming or not) and (a) time to smoking, (b) smoking duration, (c) time to incandescence and (d) incandescence duration. Species' values are within shaded probability densities, and have been jittered on the y-axis to increase visibility. Where there is a significant relationship, density shapes are coloured blue for the absence of flaming, and red for the presence of flaming. 50

Fig. 3.4 Relationship between the ignitibility index and sustainability index for the 60 study species. High y-axis values are represented by slow-ignitibility species and high x-axis values are represented by long-sustainability species. Low-combustibility species are represented by circles and high-combustibility species are represented by rhombuses. Labelled are the five low-leaf-flammability species (blue points indicating low OLFI scores) and the five high-leaf-flammability species (red points indicating high OLFI scores). 52

Fig. 4.1 Representative, fully expanded adult leaves of each of the ten study species. 62

Fig. 4.2 Probability density plots of species mean values for (a) TTI, (b) ID, (c) TTF, (d) FD and (e) the proportion of all samples flaming as a function of radiant energy. In (a) to (d), blue shaded areas show probability density, overlaid with boxplots. Letters indicate where significant differences ($P < 0.05$, after correction for multiple comparisons) exist among levels of radiant energy. Points at 29.6

kWm⁻² in (c) and (d) indicate the mean value for *B. serrata*, the only species which consistently flamed at this level of radiant energy. 67

Fig. 4.3 Mean species' values for (a) TTI, (b) ID, (c) TTF and (d) FD by radiant energy level (\pm SD). Point colours correspond to the colour key shown on the x-axis of (e). In (e), the proportion of samples flaming for each species at each level of radiant energy is shown, with light blue for 29.6 kWm⁻², red for 45.7 kWm⁻², orange for 67.6 kWm⁻² and yellow for 96.6 kWm⁻². Crosses indicate no samples flaming for a species at a given level of radiant energy. 68

Fig. 4.4 Plots of MPD (\pm SD) for (a) TTI, (b) ID, (c) TTF and (d) FD as a function of radiant energy. Letters indicate where significant differences ($P < 0.05$, after correction for multiple comparisons) exist among levels of radiant energy. In (c) and (d), letters indicate differences within two groups; species that flamed consistently from 45.7 kWm⁻² and higher, (indicated by circular points) and species that flamed consistently from 67.6 kWm⁻² and higher (indicated by rhomboid points). Species that flamed only at 96.6 kWm⁻² were excluded. 70

Fig. 5.1 The relationship between air temperature of the muffle furnace and radiant energy derived from the equation $E = \varepsilon \cdot \sigma \cdot T^4$ 79

Fig. 5.2 Flammability attributes (a) TTI and (b) ID for the 10 study species used in the predictive modelling across the four experimental levels of radiant energy. The grey boxplots are each linked to blue shaded areas which show probability densities. 81

Fig. 5.3 Classification accuracy for non-flaming (blue shaded probability densities and boxplots), and flaming (red shaded probability densities and boxplots). In (a) the 0.5 probability value is shown as a broken line, with shaded blue and red areas indicating correct predictions; (b) shows the non-linear model predicted values for Pflaming as a broken line. In both plots, the non-flaming samples below the broken line, and flaming samples above the broken line are correctly predicted. The line in (b) can be seen to pass equidistant between the medians of the flaming and non-flaming samples. 83

Fig. 5.4 Flammability attributes compared between the predictive and validation datasets. Predictive dataset values for TTI, ID and flaming behaviour are shown in (a), (b) and (c) and validation dataset

values in (b), (d) and (f). For both datasets, TTI and ID are shown as boxplots with density distributions as shaded areas. Proportions of samples flaming are indicated by shaded bars in (e, predictive) and (f, validation). In (b), (d) and (f) the species included and those not included in the predictive dataset are presented as adjacent boxplots and density distributions for each level of radiant energy (colour key shown in [b])..... 86

Fig. 5.5 (following page) Comparing mean absolute error ($MAE \pm SE$, Pearson correlation coefficient) between TTI models with (a) radiant energy as the sole explanatory variable (REM) and (b) radiant energy in conjunction with leaf traits (FM) as explanatory variables. Point colours correspond to levels of radiant energy, with a colour key shown on the x-axis of (d). The distribution of absolute error values across all observations are shown for the REM in blue and for the FM in red in (c), and for each model at each level of radiant energy in (d). Data are presented as boxplots and associated probability distributions (shaded areas). Coloured lines in (d) are the absolute error and model x radiant energy coefficients from the generalised linear model (Poisson distribution)..... 89

Fig. 5.6 (previous page) Comparing mean absolute error ($MAE \pm SE$, Pearson correlation coefficient) between ID models with (a) radiant energy as the sole explanatory variable (REM) and (b) radiant energy in conjunction with leaf traits (FM) as explanatory variables. Point colours correspond to levels of radiant energy, with a colour key shown on the x-axis of (d). The distribution of absolute error values across all observations are shown for the REM in blue and for the FM in red in (c), and for each model at each level of radiant energy in (d). Data are presented as boxplots and associated probability distributions (shaded areas). Coloured lines in (d) are the absolute error and model x radiant energy coefficients from the generalised linear model (Poisson distribution)..... 93

Fig. 5.7 Barcharts of classification accuracy of the (a) REM, (b) the FM, and (c) the FLM with the application of the Pflaming approach. Bars to the left of the x-axis divisions show the counts of non-flaming samples, and flaming samples to the right. The blue and red shaded sections of these bars indicate the counts of samples which were correctly predicted (also shown as numeric proportions above or within bars)..... 94

Fig. 5.8 (following page) Plots of observed values against predicted values of TTI for the (a) predictive dataset, (b) species in the validation dataset included in the predictive dataset, and (c) species in the validation dataset not included in the predictive dataset. Points in (a) to (c) are coloured by radiant energy level, with a key shown at the right of (c). Absolute error values across all observations for each dataset's predictions are shown in (d), with red letters indicating significant differences among the groups. Absolute error values within each radiant energy level and dataset are shown in (e). Data are presented as boxplots with shaded density distributions in (d) and (e). 95

Fig. 5.9 Absolute error values of TTI for each of the three datasets shown as separate probability density distributions for each level of radiant energy. Lines are the TTI absolute error by dataset x radiant energy coefficient from the generalised linear model (Poisson). A colour key to the datasets is shown at top right. 97

Fig. 5.10 (following page) Plots of observed values against predicted values of ID for the (a) predictive dataset, (b) species in the validation dataset included in the predictive dataset, and (c) species in the validation dataset not included in the predictive dataset. Points in (a) to (c) are coloured by radiant energy level, with a key shown at the right of (c). Absolute error values across all observations for each dataset's predictions are shown in (d), with red letters indicating significant differences among the groups. Absolute error values within each radiant energy level and dataset are shown in (e). Data are presented as boxplots with shaded density distributions in (d) and (e).... 98

Fig. 5.11 Absolute error values of ID for each of the three datasets shown as separate probability density distributions for each level of radiant energy. Lines are the ID absolute error by dataset x radiant energy coefficient from the generalised linear model (Poisson). A colour key to the datasets is shown at top right. 100

Fig. 5.12 (a) Overall accuracy of predictions for leaf flaming (or not) for the three datasets and (b) classification accuracy for samples which flamed and did not by dataset. In (b), bars to the left of the x-axis divisions show the proportion of non-flaming samples, and bars to the right show the proportion of flaming samples. The blue and red shaded sections of these bars indicate the

proportions of samples which were correctly predicted. Numbers above or within bars indicate the proportion of samples correctly predicted. 102

Fig. 5.13 Classification accuracy by radiant energy level for the predictive dataset (a), the species included (b), and the species not included in the predictive dataset (c). Bars to the left of the x-axis divisions show the counts of non-flaming samples, and flaming samples to the right. The blue and red shaded sections of these bars indicate the counts of samples which were correctly predicted. Numbers above or within bars indicate the proportion of samples correctly predicted. 103

List of Tables

Table 1.1 Results from eight studies investigating the flammability of fresh leaves. The responses measured are time to incandescence (TTI), time to flaming (TTF), incandescence duration (ID) and flaming duration (FD). The number of species included in the study, the range of values in seconds, the focal study region, and study references are provided. An 'X' in the column 'Sig.' denotes a statistical test of differences in the response value among species was made (all tests were significant at $\alpha = 0.05$).....	2
Table 2.1 Results from GLMs and PGLS models (with PGLS OU models in parentheses) of flammability attributes as a function of habitat, stratum, habitat and stratum interaction, and leaf traits (leaf area, LMA, FMC).	31
Table 3.1. Multiple regression analyses relating leaf flammability indices, the Overall Leaf Flammability Index (OLFI) and leaf combustibility to the leaf traits as predictors. Abbreviations: LMA, leaf mass per area; LA, leaf area; LWC, leaf water content.....	53
Table 5.1 Mean values of flammability attributes and leaf traits (SE in parentheses) of the study species used in the predictive modelling. The flammability attributes were all measured at 700°C. For 'Flames', a 'Y' or 'N' indicates that the species either entered or didn't enter flaming combustion respectively.....	77

List of Abbreviations in this Thesis

Context	Abbreviation	Definition
Flammability attributes	TTS	Time to smoke
	SD	Smoking duration
	TTI	Time to incandescence (equivalent to time to ignition used in Chapter 2)
	ID (BD)	Incandescence duration (equivalent to burn duration used in Chapter 2)
	TTF	Time to flaming
	FD	Flaming duration
	MLR	Mass loss rate
Leaf traits	LA	Leaf area
	LMA	Leaf mass per area
	LWC	Leaf water content
	FMC	Fuel moisture content
	FLM	Fresh leaf mass
	DLM	Dry leaf mass
Miscellaneous	PGLS	Phylogenetic generalised least squares
	OU	Ornstein-Uhlenbeck
	WUI	Wildland-urban interface
	OLFI	Overall leaf flammability index
	MPD	Mean pairwise distance
	FM	Full model

REM

Radiant energy model

LFM

Leaf flaming model

Abstract

Exacerbated by global change, wildfires are causing unprecedented impacts on ecosystems, human lives and infrastructure. Thus, there is a pressing need to better understand the drivers of wildfire in this changing world. I investigated interspecific patterns in plant leaf flammability in fire-prone vegetation of eastern Australia to determine the flammability dynamics of leaves given their essential role as fuels for wildfires.

At a landscape scale, I found that leaves of forest gully species had faster ignitibility and higher combustibility relative to woodland ridgetop species. These differences were driven by strong relationships between high flammability and both large leaf area and low leaf mass per area. For these plant communities, I explored relationships among leaf ignitibility, sustainability and combustibility and showed that faster ignitibility (higher flammability) was associated with short sustainability (lower flammability), and long sustainability (higher flammability) with higher combustibility (high flammability). Given the opposing relationship between ignitibility and sustainability, I established an overall leaf flammability index to assist in identifying low-leaf-flammability species at the wildland-urban interface. Close inspection of the flammability of ten species showed that increasing radiant energy, representing increases from low to moderate intensity wildfires, led to faster ignitability and a higher proportion of leaves flaming, such that species identity became much less important for understanding flammability dynamics. However, species identity remained particularly important for leaf sustainability and time to flaming, signalling that species differences in these flammability attributes may be helpful in understanding the dynamics of moderate intensity wildfires. Finally, I built predictive models of leaf ignitibility, sustainability and combustibility using a combination of radiant energy and leaf area, mass per area, and water content, and tested the accuracy of the models using a validation

dataset. Including leaf traits in these models dramatically increased the ability to predict leaf flammability. One application of these models will be to estimate leaf flammability for large numbers of plant species using just their leaf traits, without the need for flammability experiments if they are not feasible.

This thesis provides novel insight into interspecific variation in plant leaf flammability. The findings of my research advance our ecological understanding of leaf flammability, with potential applications to wildfire modelling, the selection of low-leaf-flammability species at the wildland-urban interface, and prediction of leaf flammability for large numbers of species using leaf traits.